



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

# ON REDUCIBLE GROUPS\*

BY

SAUL EPSTEEN

In the Transactions of the American Mathematical Society, January, 1903, pages 46–56 Professor LOEWY proves for the first time the theorem: “*Wie auch immer eine Gruppe  $G$  linearer homogener Substitutionen unter Hervorhebung ihrer irreduciblen Bestandteile in eine ähnliche Gruppe transformiert wird, so kann man die irreduciblen Bestandteilen, die sich bei irgend einer Darstellung ergeben, den irreduciblen Bestandteilen die sich bei irgend einer anderen Darstellung ergeben, eineindeutig so zuordnen, dass zwei zugeordnete irreducible Teilgruppen gleich viele Variablen haben und ähnliche Gruppen sind.*”

This says that if by transformation  $G$  be decomposed into

$$(1) \quad \begin{array}{ccccccc} \alpha_{11} & 0 & 0 & \cdots & 0 & & \\ \alpha_{21} & \alpha_{22} & 0 & \cdots & 0 & & \\ . & . & . & . & . & . & . \\ \alpha_{\lambda 1} & \alpha_{\lambda 2} & \alpha_{\lambda 3} & \cdots & \alpha_{\lambda \lambda} & & \end{array}$$

and

$$(2) \quad \begin{array}{ccccccc} \beta_{11} & 0 & 0 & \cdots & 0 & & \\ \beta_{21} & \beta_{22} & 0 & \cdots & 0 & & \\ . & . & . & . & . & . & . \\ \beta_{\mu 1} & \beta_{\mu 2} & \beta_{\mu 3} & \cdots & \beta_{\mu \mu} & & \end{array}$$

the irreducible groups

$$\begin{array}{cccc} \alpha_{11} & \alpha_{22} & \cdots & \alpha_{\lambda \lambda} \\ \beta_{11} & \beta_{22} & \cdots & \beta_{\mu \mu} \end{array}$$

are similar to each other in *some* order, that the similar group  $\alpha_{ii}$  and  $\beta_{jj}$  have the same number of variables, and that  $\lambda = \mu$ .

The object of this note is to prove the same theorem more briefly. It is well to notice however that LOEWY's proof has the advantage of being direct, while mine, although much shorter, is indirect.

Suppose we have before us a reducible linear homogeneous group  $G$  in  $n$  variables, then there exists a system of linear differential equations which have

\* Presented to the Society February 28, 1903. Received for publication February 21, 1903.

$G$  for their group of rationality.\* These equations are all of the same order and of the same type.† Let  $P_n y = 0$  be some one of these linear differential equations whose group of rationality is  $G$ . Since by hypothesis  $G$  is reducible,  $P_n y = 0$  is also reducible.‡ We now decompose  $P_n$  into irreducible factors; this may of course be accomplished in various ways:

$$(3) \quad P_n = Q_\lambda Q_{\lambda-1} \cdots Q_2 Q_1 = 0,$$

$$(4) \quad P_n = R_\mu R_{\mu-1} \cdots R_2 R_1 = 0.$$

To these decompositions of the equation there are corresponding decompositions of  $G$ , namely (1) and (2), where  $\alpha_{ii}$  is the group of  $Q_i = 0$  ( $i = 1 \cdots \lambda$ ) and  $\beta_{jj}$  is the group of  $R_j = 0$  ( $j = 1 \cdots \mu$ ).

Now LOEWY showed in his paper entitled *Ueber reduzible lineare homogene Differentialgleichungen*§ for the decompositions (3) and (4) that  $\lambda = \mu$ , and that, for suitable values of  $i$  and  $j$ ,  $Q_i$  and  $R_j$  are of the same order and type.

Since  $\lambda = \mu$  it follows that the number of irreducible groups  $\alpha$  equals the number of irreducible groups  $\beta$ . Since  $Q_i$  and  $R_j$  are of the same order, the number of variables in  $\alpha_{ii}$  is the same as the number of variables in  $\beta_{jj}$  (the number of variables in the group being the same as the order of the differential equation). Finally, since  $Q_i$  and  $R_j$  are of the same type,  $\alpha_{ii}$  and  $\beta_{jj}$  must be similar.||

The theorem may be stated in the following manner:\*\* *Considering similar groups as equivalent, the irreducible constituents of a linear homogeneous group  $G$  which are obtained by the various possible decompositions, are equivalent apart from the arrangement.*

THE UNIVERSITY OF CHICAGO.

February 20, 1903.

\* JORDAN, Bulletin de la Société Mathématique de France, vol. 2, p. 104; KLEIN, *Höhere Geometrie*, II, pp. 296-7; LOEWY, *Mathematische Annalen*, January, 1903, p. 560. For this existence theorem, see SCHLESINGER, *Handbuch* II, 1, pp. 76-7, 93.

† The French *espèce* might be translated *species*, and the German *Art* as *kind*. But in LOEWY's paper in the *Annalen* of January, 1903, p. 560, it appears that the groups of these equations are all of the same type. For this reason I say that the equations are of the same type. In accordance with the customary practice, LOEWY calls such groups equivalent.

‡ SCHLESINGER, *Handbuch*, II, 1, p. 106.

§ *Mathematische Annalen*, January, 1903, p. 565.

|| According to SCHLESINGER's *Handbuch*, II, 1, p. 121,  $\alpha_{ii}$  and  $\beta_{jj}$  are the same; but SCHLESINGER means by this that they are of the same type, i. e., can be transformed into each other. In this connection, cf. VESSIOT, *Annales de l'École Normale Supérieure*, 1892, p. 232.

\*\* Cf. LOEWY, *Transactions*, January, 1903, p. 47.